

Librería
Bonilla y Asociados
desde 1950



Título:

Autor:

Precio: \$632.00

Editorial:

Año: 2008

Tema:

Edición: 1ª

Sinopsis

ISBN: 9783540783787

In this volume the author further develops his philosophy of quantum interpolation between the real numbers and the p-adic numbers. The p-adic numbers contain the p-adic integers \mathbb{Z}_p which are the inverse limit of the finite rings \mathbb{Z}/p^n . This gives rise to a tree, and probability measures w on \mathbb{Z}_p correspond to Markov chains on this tree. From the tree structure one obtains special basis for the Hilbert space $L^2(\mathbb{Z}_p, w)$. The real analogue of the p-adic integers is the interval $[-1, 1]$, and a probability measure w on it gives rise to a special basis for $L^2([-1, 1], w)$ - the orthogonal polynomials, and to a Markov chain on "finite approximations" of $[-1, 1]$. For special (gamma and beta) measures there is a "quantum" or "q-analogue" Markov chain, and a special basis, that within certain limits yield the real and the p-adic theories. This idea can be generalized variously. In representation theory, it is the quantum general linear group $GL_n(q)$ that interpolates between the p-adic group $GL_n(\mathbb{Z}_p)$, and between its real (and complex) analogue - the orthogonal O_n (and unitary U_n) groups. There is a similar quantum interpolation between the real and p-adic Fourier transform and between the real and p-adic (local unramified part of) Tate thesis, and Weil explicit sums.