

Librería
Bonilla y Asociados
desde 1950



Título:

Autor:

Precio: \$836.00

Editorial:

Año: 2009

Tema:

Edición: 1ª

Sinopsis

ISBN: 9780821844625

Consider representation theory associated to symmetric groups, or to Hecke algebras in type A, or to q -Schur algebras, or to finite general linear groups in non-describing characteristic. Rock blocks are certain combinatorially defined blocks appearing in such a representation theory, first observed by R. Rouquier. Rock blocks are much more symmetric than general blocks, and every block is derived equivalent to a Rock block. Motivated by a theorem of J. Chuang and R. Kessar in the case of symmetric group blocks of abelian defect, the author pursues a structure theorem for these blocks.

Table of Contents

Introduction

Highest weight categories, q -Schur algebras, Hecke algebras, and finite general linear groups

Blocks of q -Schur algebras, Hecke algebras, and finite general linear groups

Rock blocks of finite general linear groups and Hecke algebras, when $w < l$

Rock blocks of symmetric groups, and the Brauer morphism

Schur-Weyl duality inside Rock blocks of symmetric groups

Ringel duality inside Rock blocks of symmetric groups

James adjustment algebras for Rock blocks of symmetric groups

Doubles, Schur super-bialgebras, and Rock blocks of Hecke algebras

Power sums

Schur doubles of type A_{∞}

Bibliography

Index